

ECE276A: Sensing & Estimation in Robotics

Lecture 10: Kalman Filtering

Instructor:

Nikolay Atanasov: nataanasov@ucsd.edu

Teaching Assistants:

Mo Shan: moshan@eng.ucsd.edu

Arash Asgharivaskasi: aasghari@eng.ucsd.edu



Bayes Filter

► **Markov Assumptions:**

- **Motion model:** given \mathbf{x}_t , \mathbf{u}_t , the state \mathbf{x}_{t+1} is independent of the history $\mathbf{x}_{0:t-1}$, $\mathbf{z}_{0:t-1}$, $\mathbf{u}_{0:t-1}$:

$$\mathbf{x}_{t+1} = f(\mathbf{x}_t, \mathbf{u}_t, \mathbf{w}_t) \sim p_f(\cdot \mid \mathbf{x}_t, \mathbf{u}_t)$$

- **Observation model:** given \mathbf{x}_t , the observation \mathbf{z}_t is independent of the history $\mathbf{x}_{0:t-1}$, $\mathbf{z}_{0:t-1}$, $\mathbf{u}_{0:t-1}$:

$$\mathbf{z}_t = h(\mathbf{x}_t, \mathbf{v}_t) \sim p_h(\cdot \mid \mathbf{x}_t)$$

- **Prior:** $p_{t|t}(\mathbf{x}_t) := p(\mathbf{x}_t \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t-1})$

- **Prediction:** $p_{t+1|t}(\mathbf{x}) = \int p_f(\mathbf{x} \mid \mathbf{s}, \mathbf{u}_t) p_{t|t}(\mathbf{s}) d\mathbf{s}$

- **Update:** $p_{t+1|t+1}(\mathbf{x}) = \frac{p_h(\mathbf{z}_{t+1} \mid \mathbf{x}) p_{t+1|t}(\mathbf{x})}{\int p_h(\mathbf{z}_{t+1} \mid \mathbf{s}) p_{t+1|t}(\mathbf{s}) d\mathbf{s}}$

Kalman Filter

- ▶ A Bayes filter with the following **assumptions**:
 - ▶ The prior pdf $p_{t|t}$ is Gaussian
 - ▶ The motion model is linear in the state \mathbf{x}_t with Gaussian noise \mathbf{w}_t
 - ▶ The observation model is linear in the state \mathbf{x}_t with Gaussian noise \mathbf{v}_t
 - ▶ The motion noise \mathbf{w}_t and observation noise \mathbf{v}_t are independent of each other, of the state \mathbf{x}_t , and across time

- ▶ **Prior:** $\mathbf{x}_t \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t-1} \sim \mathcal{N}(\boldsymbol{\mu}_{t|t}, \boldsymbol{\Sigma}_{t|t})$

- ▶ **Motion Model:**

$$\mathbf{x}_{t+1} = f(\mathbf{x}_t, \mathbf{u}_t, \mathbf{w}_t) := F\mathbf{x}_t + G\mathbf{u}_t + \mathbf{w}_t, \quad \mathbf{w}_t \sim \mathcal{N}(0, W)$$

$$\mathbf{x}_{t+1} \mid \mathbf{x}_t, \mathbf{u}_t \sim \mathcal{N}(F\mathbf{x}_t + G\mathbf{u}_t, W), \quad F \in \mathbb{R}^{d_x \times d_x}, G \in \mathbb{R}^{d_x \times d_u}, W \in \mathbb{R}^{d_x \times d_x}$$

- ▶ **Observation Model:**

$$\mathbf{z}_t = h(\mathbf{x}_t, \mathbf{v}_t) := H\mathbf{x}_t + \mathbf{v}_t, \quad \mathbf{v}_t \sim \mathcal{N}(0, V)$$

$$\mathbf{z}_t \mid \mathbf{x}_t \sim \mathcal{N}(H\mathbf{x}_t, V), \quad H \in \mathbb{R}^{d_z \times d_x}, V \in \mathbb{R}^{d_z \times d_z}$$

Matrix Manipulation

- The following results are necessary for deriving the Kalman filter:
 - **Matrix inversion lemma:**

$$(A + BDC)^{-1} = A^{-1} - A^{-1}B(D^{-1} + CA^{-1}B)^{-1}CA^{-1}$$

- **Matrix block inversion:**

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} (A - BD^{-1}C)^{-1} & -(A - BD^{-1}C)^{-1}BD^{-1} \\ -D^{-1}C(A - BD^{-1}C)^{-1} & \underbrace{D^{-1} + D^{-1}C(A - BD^{-1}C)^{-1}BD^{-1}}_{(D - CA^{-1}B)^{-1}} \end{bmatrix}$$

- **Schur complement:** $\det\left(\begin{bmatrix} A & B \\ C & D \end{bmatrix}\right) = \det(A)\det(S_A) = \det(D)\det(S_D)$

- $S_A = D - CA^{-1}B$
- $S_D = A - BD^{-1}C$

- **Square completion:**

$$\frac{1}{2}\mathbf{x}^\top A\mathbf{x} + \mathbf{b}^\top \mathbf{x} + \mathbf{c} = \frac{1}{2}(\mathbf{x} + A^{-1}\mathbf{b})^\top A(\mathbf{x} + A^{-1}\mathbf{b}) - \frac{1}{2}\mathbf{b}^\top A^{-1}\mathbf{b} + \mathbf{c}$$

Gaussian Distribution

- ▶ **Gaussian random vector** $X \sim \mathcal{N}(\mu, \Sigma)$
 - ▶ parameters:
 - ▶ **mean** $\mu \in \mathbb{R}^d$
 - ▶ **covariance** $\Sigma \in \mathbb{S}_{\succ 0}^d$ (symmetric positive definite matrix)
 - ▶ pdf: $\phi(\mathbf{x}; \mu, \Sigma) := \frac{1}{\sqrt{(2\pi)^d \det(\Sigma)}} \exp\left(-\frac{1}{2}(\mathbf{x} - \mu)^\top \Sigma^{-1}(\mathbf{x} - \mu)\right)$
 - ▶ expectation: $\mathbb{E}[X] = \int \mathbf{x} \phi(\mathbf{x}; \mu, \Sigma) d\mathbf{x} = \mu$
 - ▶ (co)variance: $\text{Var}[X] = \mathbb{E}\left[(X - \mathbb{E}[X])(X - \mathbb{E}[X])^\top\right] = \Sigma$

Information Form of the Gaussian Distribution

- ▶ An alternative Gaussian parametrization: $X \sim \mathcal{G}(\boldsymbol{\nu}, \Omega)$

- ▶ parameters:

- ▶ **information mean** $\boldsymbol{\nu} := \Sigma^{-1}\boldsymbol{\mu} \in \mathbb{R}^d$
- ▶ **information matrix**: $\Omega := \Sigma^{-1} \in \mathbb{S}_{>0}^d$

- ▶ pdf: obtained using square completion:

$$\begin{aligned}\phi(\mathbf{x}; \boldsymbol{\nu}, \Omega) &= \sqrt{\frac{\det(\Omega)}{(2\pi)^d}} \exp\left(-\frac{1}{2} (\mathbf{x}^\top \Omega \mathbf{x} - 2\boldsymbol{\nu}^\top \mathbf{x} + \boldsymbol{\nu}^\top \Omega^{-1} \boldsymbol{\nu})\right) \\ &\propto \exp\left(-\frac{1}{2} \mathbf{x}^\top \Omega \mathbf{x} + \boldsymbol{\nu}^\top \mathbf{x}\right)\end{aligned}$$

- ▶ expectation: $\mathbb{E}[X] = \Omega^{-1}\boldsymbol{\nu}$
- ▶ (co)variance: $Var[X] = \Omega^{-1}$

Covariance and Information Matrix Relationship

- ▶ The matrix block inversion relates the elements of Ω and Σ
- ▶ Let $\Omega = \begin{bmatrix} \Omega_{AA} & \Omega_{AB} \\ \Omega_{AB}^\top & \Omega_{BB} \end{bmatrix} = \Sigma^{-1} = \begin{bmatrix} \Sigma_{AA} & \Sigma_{AB} \\ \Sigma_{AB}^\top & \Sigma_{BB} \end{bmatrix}^{-1}$
- ▶ The blocks are related via:

$$\Omega_{AA} = (\Sigma_{AA} - \Sigma_{AB}\Sigma_{BB}^{-1}\Sigma_{AB}^\top)^{-1}$$

$$\Omega_{AB} = -\Omega_{AA}\Sigma_{AB}\Sigma_{BB}^{-1}$$

$$\begin{aligned}\Omega_{BB} &= \Sigma_{BB}^{-1} + \Sigma_{BB}^{-1}\Sigma_{AB}^\top\Omega_{AA}\Sigma_{AB}\Sigma_{BB}^{-1} \\ &= (\Sigma_{BB} - \Sigma_{AB}^\top\Sigma_{AA}^{-1}\Sigma_{AB})^{-1}\end{aligned}$$

- ▶ The determinants are related via:

$$\det(\Sigma) = \det(\Sigma_{AA})\det(\Omega_{BB}^{-1}) = \det(\Sigma_{BB})\det(\Omega_{AA}^{-1})$$

$$\det(\Omega) = \det(\Omega_{AA})\det(\Sigma_{BB}^{-1}) = \det(\Omega_{BB})\det(\Sigma_{AA}^{-1})$$

Gaussian Marginals and Conditionals

- ▶ Consider a **joint Gaussian distribution**:

$$\mathbf{x} := \begin{pmatrix} \mathbf{x}_A \\ \mathbf{x}_B \end{pmatrix} \sim \mathcal{N}\left(\underbrace{\begin{pmatrix} \boldsymbol{\mu}_A \\ \boldsymbol{\mu}_B \end{pmatrix}}_{\boldsymbol{\mu}}, \underbrace{\begin{bmatrix} \boldsymbol{\Sigma}_{AA} & \boldsymbol{\Sigma}_{AB} \\ \boldsymbol{\Sigma}_{AB}^\top & \boldsymbol{\Sigma}_{BB} \end{bmatrix}}_{\boldsymbol{\Sigma}}\right)$$

- ▶ The **marginal distributions** are also Gaussian:

$$\mathbf{x}_A \sim \mathcal{N}(\boldsymbol{\mu}_A, \boldsymbol{\Sigma}_{AA}) \quad \mathbf{x}_B \sim \mathcal{N}(\boldsymbol{\mu}_B, \boldsymbol{\Sigma}_{BB})$$

- ▶ The **conditional distributions** are also Gaussian:

$$\mathbf{x}_B | \mathbf{x}_A \sim \mathcal{N}\left(\boldsymbol{\mu}_B + \boldsymbol{\Sigma}_{AB}^\top \boldsymbol{\Sigma}_{AA}^{-1} (\mathbf{x}_A - \boldsymbol{\mu}_A), \underbrace{\boldsymbol{\Sigma}_{BB} - \boldsymbol{\Sigma}_{AB}^\top \boldsymbol{\Sigma}_{AA}^{-1} \boldsymbol{\Sigma}_{AB}}_{\text{Schur complement of } \boldsymbol{\Sigma}_{AA}}\right)$$

$$\mathbf{x}_A | \mathbf{x}_B \sim \mathcal{N}\left(\boldsymbol{\mu}_A + \boldsymbol{\Sigma}_{AB} \boldsymbol{\Sigma}_{BB}^{-1} (\mathbf{x}_B - \boldsymbol{\mu}_B), \underbrace{\boldsymbol{\Sigma}_{AA} - \boldsymbol{\Sigma}_{AB} \boldsymbol{\Sigma}_{BB}^{-1} \boldsymbol{\Sigma}_{AB}^\top}_{\text{Schur complement of } \boldsymbol{\Sigma}_{BB}}\right)$$

Gaussian Marginal

- Let $\tilde{\mathbf{x}}_A := \mathbf{x}_A - \boldsymbol{\mu}_A \in \mathbb{R}^n$ and $\tilde{\mathbf{x}}_B := \mathbf{x}_B - \boldsymbol{\mu}_B \in \mathbb{R}^m$ and consider:

$$\begin{aligned}
 p(\mathbf{x}_A) &= \int \phi\left(\begin{pmatrix} \mathbf{x}_A \\ \mathbf{x}_B \end{pmatrix}; \begin{pmatrix} \boldsymbol{\mu}_A \\ \boldsymbol{\mu}_B \end{pmatrix}, \begin{bmatrix} \Sigma_{AA} & \Sigma_{AB} \\ \Sigma_{AB}^\top & \Sigma_{BB} \end{bmatrix}\right) d\mathbf{x}_B \\
 &= \frac{1}{(2\pi)^{\frac{n+m}{2}} \det(\Sigma)^{1/2}} \int \exp\left(-\frac{1}{2} \left(\tilde{\mathbf{x}}_A^\top \Omega_{AA} \tilde{\mathbf{x}}_A + 2\tilde{\mathbf{x}}_A^\top \Omega_{AB} \tilde{\mathbf{x}}_B + \tilde{\mathbf{x}}_B^\top \Omega_{BB} \tilde{\mathbf{x}}_B \right)\right) d\tilde{\mathbf{x}}_B \\
 &\xrightarrow{\text{Sq. Comp.}} \frac{1}{(2\pi)^{\frac{n+m}{2}} \det(\Sigma)^{1/2}} \int \exp\left(-\frac{1}{2} \left[(\tilde{\mathbf{x}}_B + \Omega_{BB}^{-1} \Omega_{AB}^\top \tilde{\mathbf{x}}_A)^\top \Omega_{BB} (\tilde{\mathbf{x}}_B + \Omega_{BB}^{-1} \Omega_{AB}^\top \tilde{\mathbf{x}}_A) \right.\right. \\
 &\quad \left.\left. - \tilde{\mathbf{x}}_A^\top \Omega_{AB} \Omega_{BB}^{-1} \Omega_{AB}^\top \tilde{\mathbf{x}}_A + \tilde{\mathbf{x}}_A^\top \Omega_{AA} \tilde{\mathbf{x}}_A \right]\right) d\tilde{\mathbf{x}}_B
 \end{aligned}$$

- Note that:

- $\int \phi(\tilde{\mathbf{x}}_B; -\Omega_{BB}^{-1} \Omega_{AB}^\top \tilde{\mathbf{x}}_A, \Omega_{BB}^{-1}) d\tilde{\mathbf{x}}_B = 1$
- $\Sigma_{AA}^{-1} = \Omega_{AA} - \Omega_{AB} \Omega_{BB}^{-1} \Omega_{AB}^\top$
- $\det(\Sigma) = \det(\Sigma_{AA}) \det(\Omega_{BB}^{-1})$

$$\begin{aligned}
 &= \frac{1}{(2\pi)^{\frac{n}{2}} \det(\Sigma_{AA})^{1/2}} \exp\left(-\frac{1}{2} \tilde{\mathbf{x}}_A^\top \Sigma_{AA}^{-1} \tilde{\mathbf{x}}_A\right) \int \phi(\tilde{\mathbf{x}}_B; -\Omega_{BB}^{-1} \Omega_{AB}^\top \tilde{\mathbf{x}}_A, \Omega_{BB}^{-1}) d\tilde{\mathbf{x}}_B
 \end{aligned}$$

↑ 1

$\Rightarrow \mathbf{x}_A \sim \mathcal{N}(\boldsymbol{\mu}_A, \Sigma_{AA})$

Gaussian Conditional

$$\begin{aligned} p(\mathbf{x}_A \mid \mathbf{x}_B) &= \frac{p(\mathbf{x}_A, \mathbf{x}_B)}{p(\mathbf{x}_B)} = \frac{\frac{1}{\sqrt{(2\pi)^{n+m} \det(\Sigma)}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^\top \Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu})}}{\frac{1}{\sqrt{(2\pi)^m \det(\Sigma_{BB})}} e^{-\frac{1}{2}(\mathbf{x}_B-\boldsymbol{\mu}_B)^\top \Sigma_{BB}^{-1}(\mathbf{x}_B-\boldsymbol{\mu}_B)}} \\ &= \frac{1}{\sqrt{(2\pi)^n \det(\Sigma) / \det(\Sigma_{BB})}} e^{-\frac{1}{2}\left((\mathbf{x}-\boldsymbol{\mu})^\top \Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu}) - (\mathbf{x}_B-\boldsymbol{\mu}_B)^\top \Sigma_{BB}^{-1}(\mathbf{x}_B-\boldsymbol{\mu}_B)\right)} \end{aligned}$$

► Consider the exponent:

$$\begin{aligned} &\tilde{\mathbf{x}}_A^\top \Omega_{AA} \tilde{\mathbf{x}}_A + \tilde{\mathbf{x}}_A^\top \Omega_{AB} \tilde{\mathbf{x}}_B + \tilde{\mathbf{x}}_B^\top \Omega_{AB}^\top \tilde{\mathbf{x}}_A + \tilde{\mathbf{x}}_B^\top \Omega_{BB} \tilde{\mathbf{x}}_B - \tilde{\mathbf{x}}_B^\top \Sigma_{BB}^{-1} \tilde{\mathbf{x}}_B \\ &= \tilde{\mathbf{x}}_A^\top \Omega_{AA} \tilde{\mathbf{x}}_A - 2\tilde{\mathbf{x}}_A^\top \Omega_{AA} \Sigma_{AB} \Sigma_{BB}^{-1} \tilde{\mathbf{x}}_B + \tilde{\mathbf{x}}_B^\top \Sigma_{BB}^{-1} \Sigma_{AB}^\top \Omega_{AA} \Sigma_{AB} \Sigma_{BB}^{-1} \tilde{\mathbf{x}}_B \\ &= (\tilde{\mathbf{x}}_A - \Sigma_{AB} \Sigma_{BB}^{-1} \tilde{\mathbf{x}}_B)^\top \Omega_{AA} (\tilde{\mathbf{x}}_A - \Sigma_{AB} \Sigma_{BB}^{-1} \tilde{\mathbf{x}}_B) \end{aligned}$$

$$\Rightarrow \boxed{\mathbf{x}_A \mid \mathbf{x}_B \sim \mathcal{N} \left(\boldsymbol{\mu}_A + \Sigma_{AB} \Sigma_{BB}^{-1} (\mathbf{x}_B - \boldsymbol{\mu}_B), \Sigma_{AA} - \Sigma_{AB} \Sigma_{BB}^{-1} \Sigma_{AB}^\top \right)}$$

The Gaussian Distribution is Stable

- ▶ **Stable distribution:** a linear combination $aX_1 + bX_2$ of two independent copies of a random variable X has the same distribution as $cX + d$ up to location d and scale $c > 0$ parameters
- ▶ The Gaussian distribution is stable
- ▶ Since addition of random variables corresponds to convolution of their pdfs, the space of Gaussian pdfs is **closed under convolution**:

$$\int \phi(\mathbf{x}; F\mathbf{s}, W) \phi(\mathbf{s}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{s} = \phi\left(\mathbf{x}; F\boldsymbol{\mu}, F\boldsymbol{\Sigma}F^\top + W\right)$$

- ▶ The space of Gaussian pdfs is also **closed under geometric averages** (up to scaling):

$$\prod_k \phi^{\alpha_k}(\mathbf{x}; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \propto \phi\left(\mathbf{x}; \left(\sum_k \alpha_k \boldsymbol{\Sigma}_k^{-1}\right)^{-1} \left(\sum_k \boldsymbol{\Sigma}_k^{-1} \boldsymbol{\mu}_k\right), \left(\sum_k \alpha_k \boldsymbol{\Sigma}_k^{-1}\right)^{-1}\right)$$

Kalman Filter

- ▶ A Bayes filter with the following **assumptions**:
 - ▶ The prior pdf $p_{t|t}$ is Gaussian
 - ▶ The motion model is linear in the state \mathbf{x}_t with Gaussian noise \mathbf{w}_t
 - ▶ The observation model is linear in the state \mathbf{x}_t with Gaussian noise \mathbf{v}_t
 - ▶ The motion noise \mathbf{w}_t and observation noise \mathbf{v}_t are independent of each other, of the state \mathbf{x}_t , and across time

- ▶ **Prior:** $\mathbf{x}_t | \mathbf{z}_{0:t}, \mathbf{u}_{0:t-1} \sim \mathcal{N}(\boldsymbol{\mu}_{t|t}, \boldsymbol{\Sigma}_{t|t})$

- ▶ **Motion Model:**

$$\mathbf{x}_{t+1} = f(\mathbf{x}_t, \mathbf{u}_t, \mathbf{w}_t) := F\mathbf{x}_t + G\mathbf{u}_t + \mathbf{w}_t, \quad \mathbf{w}_t \sim \mathcal{N}(0, W)$$

$$\mathbf{x}_{t+1} | \mathbf{x}_t, \mathbf{u}_t \sim \mathcal{N}(F\mathbf{x}_t + G\mathbf{u}_t, W), \quad F \in \mathbb{R}^{d_x \times d_x}, G \in \mathbb{R}^{d_x \times d_u}, W \in \mathbb{R}^{d_x \times d_x}$$

- ▶ **Observation Model:**

$$\mathbf{z}_t = h(\mathbf{x}_t, \mathbf{v}_t) := H\mathbf{x}_t + \mathbf{v}_t, \quad \mathbf{v}_t \sim \mathcal{N}(0, V)$$

$$\mathbf{z}_t | \mathbf{x}_t \sim \mathcal{N}(H\mathbf{x}_t, V), \quad H \in \mathbb{R}^{d_z \times d_x}, V \in \mathbb{R}^{d_z \times d_z}$$

Kalman Filter Prediction

$$\begin{aligned} p_{t+1|t}(\mathbf{x}) &= \int p_f(\mathbf{x} \mid \mathbf{s}, \mathbf{u}_t) p_{t|t}(\mathbf{s}) d\mathbf{s} = \int \phi(\mathbf{x}; F\mathbf{s} + G\mathbf{u}_t, W) \phi(\mathbf{s}; \boldsymbol{\mu}_{t|t}, \boldsymbol{\Sigma}_{t|t}) d\mathbf{s} \\ &= \underbrace{\frac{1}{(2\pi)^{d_x} \sqrt{\det(W) \det(\boldsymbol{\Sigma}_{t|t})}}}_{\kappa_{t|t}} \int \exp \left\{ -\frac{1}{2} (\mathbf{x} - F\mathbf{s} - G\mathbf{u}_t)^\top W^{-1} (\mathbf{x} - F\mathbf{s} - G\mathbf{u}_t) \right\} \times \\ &\quad \exp \left\{ -\frac{1}{2} (\mathbf{s} - \boldsymbol{\mu}_{t|t})^\top \boldsymbol{\Sigma}_{t|t}^{-1} (\mathbf{s} - \boldsymbol{\mu}_{t|t}) \right\} d\mathbf{s} \\ &= \kappa_{t|t} \int \exp \left\{ -\frac{1}{2} \left(\mathbf{s}^\top (F^\top W^{-1} F + \boldsymbol{\Sigma}_{t|t}^{-1}) \mathbf{s} - 2(\boldsymbol{\Sigma}_{t|t}^{-1} \boldsymbol{\mu}_{t|t} + F^\top W^{-1} (\mathbf{x} - G\mathbf{u}_t))^\top \mathbf{s} + \dots \right) \right\} d\mathbf{s} \\ &\stackrel{\substack{\text{Sq. Comp.} \\ \text{Inv. Lemma}}}{=} \phi(\mathbf{x}; F\boldsymbol{\mu}_{t|t} + G\mathbf{u}_t, F\boldsymbol{\Sigma}_{t|t}F^\top + W) \end{aligned}$$

$$\boxed{\begin{aligned} p_{t+1|t}(\mathbf{x}) &= \int \phi(\mathbf{x}; F\mathbf{s} + G\mathbf{u}_t, W) \phi(\mathbf{s}; \boldsymbol{\mu}_{t|t}, \boldsymbol{\Sigma}_{t|t}) d\mathbf{s} \\ &= \phi(\mathbf{x}; F\boldsymbol{\mu}_{t|t} + G\mathbf{u}_t, F\boldsymbol{\Sigma}_{t|t}F^\top + W) \end{aligned}}$$

Kalman Filter Prediction (easy version)

- ▶ Motion model with given prior:

$$\mathbf{x}_{t+1} = F\mathbf{x}_t + G\mathbf{u}_t + \mathbf{w}_t, \quad \mathbf{w}_t \sim \mathcal{N}(0, W), \quad \mathbf{x}_t \sim \mathcal{N}(\boldsymbol{\mu}_{t|t}, \Sigma_{t|t})$$

- ▶ Since \mathbf{w}_t and \mathbf{x}_t are independent and the Gaussian distribution is stable, we know that the distribution of \mathbf{x}_{t+1} is Gaussian: $\mathcal{N}(\boldsymbol{\mu}_{t+1|t}, \Sigma_{t+1|t})$
- ▶ We just need to compute its mean and covariance:

$$\boldsymbol{\mu}_{t+1|t} = \mathbb{E}[F\mathbf{x}_t + G\mathbf{u}_t + \mathbf{w}_t] = F\mathbb{E}[\mathbf{x}_t] + G\mathbf{u}_t + \mathbb{E}[\mathbf{w}_t] = F\boldsymbol{\mu}_{t|t} + G\mathbf{u}_t$$

$$\begin{aligned}\Sigma_{t+1|t} &= \mathbf{Var}[F\mathbf{x}_t + G\mathbf{u}_t + \mathbf{w}_t] \xrightarrow{\text{independence}} \mathbf{Var}[F\mathbf{x}_t] + 0 + \mathbf{Var}[\mathbf{w}_t] \\ &= \mathbb{E}\left[F(\mathbf{x}_t - \boldsymbol{\mu}_{t|t})(\mathbf{x}_t - \boldsymbol{\mu}_{t|t})^\top F^\top\right] + W \\ &= F\Sigma_{t|t}F^\top + W\end{aligned}$$

Kalman Filter Update

$$\begin{aligned} p_{t+1|t+1}(\mathbf{x}) &= \frac{p(\mathbf{z}_{t+1} | \mathbf{x}) p_{t+1|t}(\mathbf{x})}{p(\mathbf{z}_{t+1} | \mathbf{z}_{0:t}, \mathbf{u}_{0:t})} = \frac{\phi(\mathbf{z}_{t+1}; H\mathbf{x}, V) \phi(\mathbf{x}; \boldsymbol{\mu}_{t+1|t}, \boldsymbol{\Sigma}_{t+1|t})}{\int \phi(\mathbf{z}_{t+1}; H\mathbf{s}, V) \phi(\mathbf{s}; \boldsymbol{\mu}_{t+1|t}, \boldsymbol{\Sigma}_{t+1|t}) d\mathbf{s}} \\ &= \frac{\phi(\mathbf{z}_{t+1}; H\mathbf{x}, V) \phi(\mathbf{x}; \boldsymbol{\mu}_{t+1|t}, \boldsymbol{\Sigma}_{t+1|t})}{\phi(\mathbf{z}_{t+1}; H\boldsymbol{\mu}_{t+1|t}, H\boldsymbol{\Sigma}_{t+1|t}H^\top + V)} \\ &= \frac{\kappa_{t+1}}{\eta_{t+1}} \exp \left\{ -\frac{1}{2} (\mathbf{z}_{t+1} - H\mathbf{x})^\top V^{-1} (\mathbf{z}_{t+1} - H\mathbf{x}) \right\} \exp \left\{ -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu}_{t+1|t})^\top \boldsymbol{\Sigma}_{t+1|t}^{-1} (\mathbf{x} - \boldsymbol{\mu}_{t+1|t}) \right\} \\ &= \frac{\kappa_{t+1}}{\eta_{t+1}} \exp \left\{ -\frac{1}{2} \left(\mathbf{x}^\top (H^\top V^{-1} H + \boldsymbol{\Sigma}_{t+1|t}^{-1}) \mathbf{x} + 2(H^\top V^{-1} \mathbf{z}_{t+1} + \boldsymbol{\Sigma}_{t+1|t}^{-1} \boldsymbol{\mu}_{t+1|t})^\top \mathbf{x} + \dots \right) \right\} \\ &\xrightarrow{\text{Sq. Comp.}} \phi \left(\mathbf{x}; (H^\top V^{-1} H + \boldsymbol{\Sigma}_{t+1|t}^{-1})^{-1} (H^\top V^{-1} \mathbf{z}_{t+1} + \boldsymbol{\Sigma}_{t+1|t}^{-1} \boldsymbol{\mu}_{t+1|t}), (H^\top V^{-1} H + \boldsymbol{\Sigma}_{t+1|t}^{-1})^{-1} \right) \\ &\xrightarrow{\text{Inv. Lemma}} \boxed{\phi \left(\mathbf{x}; \boldsymbol{\mu}_{t+1|t} + K_{t+1|t} (\mathbf{z}_{t+1} - H\boldsymbol{\mu}_{t+1|t}), (I - K_{t+1|t} H) \boldsymbol{\Sigma}_{t+1|t} \right)} \end{aligned}$$

- **Kalman gain:** $K_{t+1|t} := \boldsymbol{\Sigma}_{t+1|t} H^\top (H\boldsymbol{\Sigma}_{t+1|t} H^\top + V)^{-1}$

Kalman Filter Update (easy version)

- ▶ Observation model with given prior:

$$\mathbf{z}_{t+1} = H\mathbf{x}_{t+1} + \mathbf{v}_{t+1}, \quad \mathbf{v}_{t+1} \sim \mathcal{N}(0, V), \quad \mathbf{x}_{t+1} \sim \mathcal{N}(\boldsymbol{\mu}_{t+1|t}, \Sigma_{t+1|t})$$

- ▶ The joint distribution of \mathbf{x}_{t+1} and \mathbf{z}_{t+1} is Gaussian:

$$\begin{pmatrix} \mathbf{x}_{t+1} \\ \mathbf{z}_{t+1} \end{pmatrix} \sim \mathcal{N} \left(\begin{pmatrix} \boldsymbol{\mu}_{t+1|t} \\ H\boldsymbol{\mu}_{t+1|t} \end{pmatrix}, \begin{bmatrix} \Sigma_{t+1|t} & C_{t+1|t} \\ C_{t+1|t}^\top & H\Sigma_{t+1|t}H^\top + V \end{bmatrix} \right)$$

$$\begin{aligned} C_{t+1|t} &= \mathbb{E} \left[(\mathbf{x}_{t+1} - \boldsymbol{\mu}_{t+1|t}) (\mathbf{z}_{t+1} - H\boldsymbol{\mu}_{t+1|t})^\top \right] \\ &= \mathbb{E} \left[(\mathbf{x}_{t+1} - \boldsymbol{\mu}_{t+1|t}) \left[(\mathbf{x}_{t+1} - \boldsymbol{\mu}_{t+1|t})^\top H^\top + \mathbf{v}_{t+1}^\top \right] \right] = \Sigma_{t+1|t} H^\top \end{aligned}$$

- ▶ The conditional distribution of $\mathbf{x}_{t+1} | \mathbf{z}_{t+1}$ is then also Gaussian:

$$\begin{aligned} \mathbf{x}_{t+1} | \mathbf{z}_{t+1} &\sim \mathcal{N} \left(\boldsymbol{\mu}_{t+1|t} + \Sigma_{t+1|t} H^\top (H\Sigma_{t+1|t} H^\top + V)^{-1} (\mathbf{z}_{t+1} - H\boldsymbol{\mu}_{t+1|t}), \right. \\ &\quad \left. \Sigma_{t+1|t} - \Sigma_{t+1|t} H^\top (H\Sigma_{t+1|t} H^\top + V)^{-1} H\Sigma_{t+1|t} \right) \end{aligned}$$

Kalman Filter Summary

Motion model: $\mathbf{x}_{t+1} = F\mathbf{x}_t + G\mathbf{u}_t + \mathbf{w}_t, \quad \mathbf{w}_t \sim \mathcal{N}(0, W)$

Observation model: $\mathbf{z}_t = H\mathbf{x}_t + \mathbf{v}_t, \quad \mathbf{v}_t \sim \mathcal{N}(0, V)$

Prior: $\mathbf{x}_t \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t-1} \sim \mathcal{N}(\boldsymbol{\mu}_{t|t}, \boldsymbol{\Sigma}_{t|t})$

Prediction:

$$\begin{aligned}\boldsymbol{\mu}_{t+1|t} &= F\boldsymbol{\mu}_{t|t} + G\mathbf{u}_t \\ \boldsymbol{\Sigma}_{t+1|t} &= F\boldsymbol{\Sigma}_{t|t}F^\top + W\end{aligned}$$


Update:

$$\begin{aligned}\boldsymbol{\mu}_{t+1|t+1} &= \boldsymbol{\mu}_{t+1|t} + K_{t+1|t}(\mathbf{z}_{t+1} - H\boldsymbol{\mu}_{t+1|t}) \\ \boldsymbol{\Sigma}_{t+1|t+1} &= (I - K_{t+1|t}H)\boldsymbol{\Sigma}_{t+1|t}\end{aligned}$$


Kalman Gain: $K_{t+1|t} := \boldsymbol{\Sigma}_{t+1|t}H^\top (H\boldsymbol{\Sigma}_{t+1|t}H^\top + V)^{-1}$

Predict


Update




Predict





Update




Predict






Update



Predict



Update



Kalman Filter Comments

- ▶ The normalization factor in the update step (Bayes rule) is Gaussian:

$$\begin{aligned} p(\mathbf{z}_{t+1} \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t}) &= \int p(\mathbf{z}_{t+1} \mid \mathbf{s}) p(\mathbf{s} \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t}) d\mathbf{s} \\ &= \int \phi(\mathbf{z}_{t+1}; H\mathbf{s}, V) \phi(\mathbf{s}; \boldsymbol{\mu}_{t+1|t}, \boldsymbol{\Sigma}_{t+1|t}) d\mathbf{s} \\ &= \phi(\mathbf{z}_{t+1}; H\boldsymbol{\mu}_{t+1|t}, H\boldsymbol{\Sigma}_{t+1|t}H^\top + V) \end{aligned}$$

- ▶ **Innovation:** the term $\mathbf{r}_{t+1} := \mathbf{z}_{t+1} - H\boldsymbol{\mu}_{t+1|t}$ used to correct the mean in the update step
- ▶ The innovation \mathbf{r}_{t+1} conditioned on the past information $\mathbf{z}_{0:t}, \mathbf{u}_{0:t}$ has the same covariance as $p(\mathbf{z}_{t+1} \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t})$:

$$\mathbb{E}[\mathbf{r}_{t+1} \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t}] = H\boldsymbol{\mu}_{t+1|t} - H\boldsymbol{\mu}_{t+1|t} = 0$$

$$\mathbb{E}[\mathbf{r}_{t+1}\mathbf{r}_{t+1}^\top \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t}] = H\boldsymbol{\Sigma}_{t+1|t}H^\top + V$$

- ▶ **Kalman gain:** the matrix $K_{t+1|t}$ scales the innovation \mathbf{r}_{t+1} by its covariance and determines how much to trust it in the update

Kalman Filter Comments

- ▶ **Efficient:** polynomial in measurement and state dim: $O(d_z^{2.376} + d_x^2)$
- ▶ **Optimal:** under linear, Gaussian, and independence assumptions with respect to the mean square error (MSE):

$$\mathbb{E} \left[\| \mathbf{x}_t - \boldsymbol{\mu}_{t|t} \|_2^2 \right] = \text{tr}(\boldsymbol{\Sigma}_{t|t})$$

- ▶ To deal with **unknown models** we can use EM to learn the motion model (F, G, W) and the observation model (H, V)
- ▶ Given data $\mathcal{D} := \{\mathbf{z}_{0:T}, \mathbf{u}_{0:T-1}\}$, apply EM with hidden variables $\mathbf{x}_{0:T}$:
 - ▶ **E step:** Given initial parameter estimates $\boldsymbol{\theta}^{(k)} := \{F^{(k)}, G^{(k)}, W^{(k)}, H^{(k)}, V^{(k)}\}$ calculate the likelihood of the hidden variables via the Kalman filter/smooth
 - ▶ **M step:** Optimize the parameters via MLE to obtain $\boldsymbol{\theta}^{(k+1)}$ which explain the posterior distribution over $\mathbf{x}_{0:T}$ better
- ▶ Most robot systems are **nonlinear!**

Information Filter

- ▶ Uses the information form $\mathbf{x} \sim \mathcal{G}(\boldsymbol{\nu}, \Omega)$ where $\boldsymbol{\nu} = \Sigma^{-1}\boldsymbol{\mu}$ and $\Omega = \Sigma^{-1}$
- ▶ Converts the Kalman filter equations to their information form counterparts via the matrix inversion lemma

Motion model: $\mathbf{x}_{t+1} = F\mathbf{x}_t + G\mathbf{u}_t + \mathbf{w}_t, \quad \mathbf{w}_t \sim \mathcal{G}(0, W^{-1})$

Observation model: $\mathbf{z}_t = H\mathbf{x}_t + \mathbf{v}_t, \quad \mathbf{v}_t \sim \mathcal{G}(0, V^{-1})$

Prior: $\mathbf{x}_t \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t-1} \sim \mathcal{G}(\boldsymbol{\nu}_{t|t}, \Omega_{t|t})$

Prediction:

$$\boldsymbol{\nu}_{t+1|t} = \Omega_{t+1|t} \left(F\Omega_{t|t}^{-1}\boldsymbol{\nu}_{t|t} + G\mathbf{u}_t \right)$$
$$\Omega_{t+1|t} = \left(F\Omega_{t|t}^{-1}F^\top + W \right)^{-1}$$

Update:

$$\boldsymbol{\nu}_{t+1|t+1} = \boldsymbol{\nu}_{t+1|t} + H^\top V^{-1} \mathbf{z}_{t+1}$$
$$\Omega_{t+1|t+1} = \Omega_{t+1|t} + H^\top V^{-1} H$$

Kalman-Bucy Filter (continuous time)

Motion model: $\dot{\mathbf{x}}(t) = F\mathbf{x}(t) + G\mathbf{u}(t) + \mathbf{w}(t)$

Observation model: $\mathbf{z}(t) = H\mathbf{x}(t) + \mathbf{v}(t)$

Prior: $\mathbf{x}(0) \sim \mathcal{N}(\mu(0), \Sigma(0))$

Mean: $\dot{\mu}(t) = F\mu(t) + G\mathbf{u}(t) + K(t)(\mathbf{z}(t) - H\mu(t))$

Covariance: $\dot{\Sigma}(t) = F\Sigma(t)F^\top + W - K(t)VK^\top(t)$

Kalman Gain: $K(t) = \Sigma(t)H^\top V^{-1}$

Gaussian Mixture Filter

► **Motion model:** $\mathbf{x}_{t+1} = F\mathbf{x}_t + G\mathbf{u}_t + \mathbf{w}_t$, $\mathbf{w}_t \sim \mathcal{N}(0, W)$

► **Observation model:** $\mathbf{z}_t = H\mathbf{x}_t + \mathbf{v}_t$, $\mathbf{v}_t \sim \mathcal{N}(0, V)$

► **Prior:** $\mathbf{x}_t | \mathbf{z}_{0:t}, \mathbf{u}_{0:t-1} \sim p_{t|t}(\mathbf{x}_t) := \sum_k \alpha_{t|t}^{(k)} \phi\left(\mathbf{x}_t; \boldsymbol{\mu}_{t|t}^{(k)}, \Sigma_{t|t}^{(k)}\right)$

► **Prediction:**

$$\begin{aligned} p_{t+1|t}(\mathbf{x}) &= \int p_f(\mathbf{x} | \mathbf{s}, \mathbf{u}_t) p_{t|t}(\mathbf{s}) d\mathbf{s} = \sum_k \alpha_{t|t}^{(k)} \int \phi(\mathbf{x}; F\mathbf{s} + G\mathbf{u}_t, W) \phi\left(\mathbf{s}; \boldsymbol{\mu}_{t|t}^{(k)}, \Sigma_{t|t}^{(k)}\right) d\mathbf{s} \\ &= \sum_k \alpha_{t|t}^{(k)} \phi\left(\mathbf{x}; F\boldsymbol{\mu}_{t|t}^{(k)} + G\mathbf{u}_t, F\Sigma_{t|t}^{(k)}F^\top + W\right) \end{aligned}$$

► **Update:**

$$\begin{aligned} p_{t+1|t+1}(\mathbf{x}) &= \frac{p_h(\mathbf{z}_{t+1} | \mathbf{x}) p_{t+1|t}(\mathbf{x})}{p(\mathbf{z}_{t+1} | \mathbf{z}_{0:t}, \mathbf{u}_{0:t})} = \frac{\phi(\mathbf{z}_{t+1}; H\mathbf{x}, V) \sum_k \alpha_{t+1|t}^{(k)} \phi\left(\mathbf{x}; \boldsymbol{\mu}_{t+1|t}^{(k)}, \Sigma_{t+1|t}^{(k)}\right)}{\int \phi(\mathbf{z}_{t+1}; H\mathbf{s}, V) \sum_j \alpha_{t+1|t}^{(j)} \phi\left(\mathbf{s}; \boldsymbol{\mu}_{t+1|t}^{(j)}, \Sigma_{t+1|t}^{(j)}\right) d\mathbf{s}} \\ &= \sum_k \left(\frac{\alpha_{t+1|t}^{(k)} \phi(\mathbf{z}_{t+1}; H\mathbf{x}, V) \phi\left(\mathbf{x}; \boldsymbol{\mu}_{t+1|t}^{(k)}, \Sigma_{t+1|t}^{(k)}\right)}{\sum_j \alpha_{t+1|t}^{(j)} \phi\left(\mathbf{z}_{t+1}; H\boldsymbol{\mu}_{t+1|t}^{(j)}, H\Sigma_{t+1|t}^{(j)}H^\top + V\right)} \times \frac{\phi(\mathbf{z}_{t+1}; H\boldsymbol{\mu}_{t+1|t}^{(k)}, H\Sigma_{t+1|t}^{(k)}H^\top + V)}{\phi(\mathbf{z}_{t+1}; H\boldsymbol{\mu}_{t+1|t}^{(k)}, H\Sigma_{t+1|t}^{(k)}H^\top + V)} \right) \\ &= \sum_k \left[\frac{\alpha_{t+1|t}^{(k)} \phi\left(\mathbf{z}_{t+1}; H\boldsymbol{\mu}_{t+1|t}^{(k)}, H\Sigma_{t+1|t}^{(k)}H^\top + V\right)}{\sum_j \alpha_{t+1|t}^{(j)} \phi\left(\mathbf{z}_{t+1}; H\boldsymbol{\mu}_{t+1|t}^{(j)}, H\Sigma_{t+1|t}^{(j)}H^\top + V\right)} \right] \phi\left(\mathbf{x}; \boldsymbol{\mu}_{t+1|t}^{(k)} + K_{t+1|t}^{(k)}(\mathbf{z}_{t+1} - H\boldsymbol{\mu}_{t+1|t}^{(k)}), (I - K_{t+1|t}^{(k)}H)\Sigma_{t+1|t}^{(k)}\right) \end{aligned}$$

► **Kalman Gain:** $K_{t+1|t}^{(k)} := \Sigma_{t+1|t}^{(k)} H^\top \left(H\Sigma_{t+1|t}^{(k)}H^\top + V \right)^{-1}$

Gaussian Mixture Filter

- ▶ The GMF is just a **bank of Kalman filters** with weights $\alpha^{(k)}$ scaled by the observation model in the update step
- ▶ The GMF is also called the **Gaussian Sum Filter**
- ▶ **pdf:** $\mathbf{x}_t \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t-1} \sim p_{t|t}(\mathbf{x}) := \sum_k \alpha_{t|t}^{(k)} \phi\left(\mathbf{x}_t; \boldsymbol{\mu}_{t|t}^{(k)}, \boldsymbol{\Sigma}_{t|t}^{(k)}\right)$
- ▶ **mean:** $\boldsymbol{\mu}_{t|t} := \mathbb{E}[\mathbf{x}_t \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t-1}] = \int \mathbf{x} p_{t|t}(\mathbf{x}) d\mathbf{x} = \sum_k \alpha_{t|t}^{(k)} \boldsymbol{\mu}_{t|t}^{(k)}$
- ▶ **covariance:** $\boldsymbol{\Sigma}_{t|t} := \mathbb{E}\left[\mathbf{x}_t \mathbf{x}_t^\top \mid \mathbf{z}_{0:t}, \mathbf{u}_{0:t-1}\right] - \boldsymbol{\mu}_{t|t} \boldsymbol{\mu}_{t|t}^\top$
 $= \int \mathbf{x} \mathbf{x}^\top p_{t|t}(\mathbf{x}) d\mathbf{x} - \boldsymbol{\mu}_{t|t} \boldsymbol{\mu}_{t|t}^\top = \sum_k \alpha_{t|t}^{(k)} \left(\boldsymbol{\Sigma}_{t|t}^{(k)} + \boldsymbol{\mu}_{t|t}^{(k)} (\boldsymbol{\mu}_{t|t}^{(k)})^\top \right) - \boldsymbol{\mu}_{t|t} \boldsymbol{\mu}_{t|t}^\top$

Rao-Blackwellized Particle Filter

- The Rao-Blackwellized (**marginalized**) particle filter is applicable to conditionally linear-Gaussian models:

$$x_{t+1}^n = f_t^n(x_t^n) + A_t^n(x_t^n)x_t^l + G_t^n(x_t^n)w_t^n$$

$$x_{t+1}^l = f_t^l(x_t^n) + A_t^l(x_t^n)x_t^l + G_t^l(x_t^n)w_t^l$$

$$z_t = h_t(x_t^n) + C_t(x_t^n)x_t^l + v_t$$

Nonlinear states: x_t^n
 Linear states: x_t^l

- Idea:** exploit linear-Gaussian sub-structure to handle high dim. problems

$$p\left(x_t^l, x_{0:t}^n \mid z_{0:t}, u_{0:t-1}\right) = \underbrace{p\left(x_t^l \mid z_{0:t}, u_{0:t-1}, x_{0:t}^n\right)}_{\text{Kalman Filter}} \underbrace{p\left(x_{0:t}^n \mid z_{0:t}, u_{0:t-1}\right)}_{\text{Particle Filter}}$$

$$= \sum_{k=1}^{N_{t|t}} \alpha_{t|t}^{(k)} \delta\left(x_{0:t}^n; m_{t|t}^{(k)}\right) \phi\left(x_t^l; \mu_{t|t}^{(k)}, \Sigma_{t|t}^{(k)}\right)$$

- The RBPF is a combination of the particle filter and the Kalman filter, in which each particle has a Kalman filter associated to it

