

ECE276A: Sensing & Estimation in Robotics

Lecture 3: Linear Algebra (Review)

Instructor:

Nikolay Atanasov: natanasov@ucsd.edu

Teaching Assistants:

Qiaojun Feng: qjfeng@ucsd.edu

Arash Asgharivaskasi: aasghari@eng.ucsd.edu

Ehsan Zobeidi: ezobeidi@ucsd.edu

Rishabh Jangir: rjangir@ucsd.edu

UC San Diego

JACOBS SCHOOL OF ENGINEERING
Electrical and Computer Engineering

Field

- ▶ A **field** is a set F with two binary operations, $+$: $F \times F \mapsto F$ (addition) and \cdot : $F \times F \mapsto F$ (multiplication), which satisfy the following axioms:
 - ▶ **Associativity**: $a + (b + c) = (a + b) + c$ and $a(bc) = (ab)c$, $\forall a, b, c \in F$
 - ▶ **Commutativity**: $a + b = b + a$ and $ab = ba$, $\forall a, b \in F$
 - ▶ **Identity**: $\exists 1, 0 \in F$ such that $a + 0 = a$ and $a1 = a$, $\forall a \in F$
 - ▶ **Inverse**: $\forall a \in F, \exists -a \in F$ such that $a + (-a) = 0$
 $\forall a \in F \setminus \{0\}, \exists a^{-1} \in F \setminus \{0\}$ such that $aa^{-1} = 1$
 - ▶ **Distributivity**: $a(b + c) = (ab) + (ac)$, $\forall a, b, c \in F$
- ▶ **Examples**: real numbers \mathbb{R} , complex numbers \mathbb{C} , rational numbers \mathbb{Q}

Vector Space

- ▶ A **vector space** over a field F is a set V with two binary operations, $+$: $V \times V \mapsto V$ (addition) and \cdot : $F \times V \mapsto V$ (scalar multiplication), which satisfy the following axioms:
 - ▶ **Associativity**: $\mathbf{x} + (\mathbf{y} + \mathbf{z}) = (\mathbf{x} + \mathbf{y}) + \mathbf{z}$, $\forall \mathbf{x}, \mathbf{y}, \mathbf{z} \in V$
 - ▶ **Compatibility**: $a(b\mathbf{x}) = (ab)\mathbf{x}$, $\forall a, b \in F$ and $\forall \mathbf{x} \in V$
 - ▶ **Commutativity**: $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$, $\forall \mathbf{x}, \mathbf{y} \in V$
 - ▶ **Identity**: $\exists \mathbf{0} \in V$ and $1 \in F$ such that $\mathbf{x} + \mathbf{0} = \mathbf{x}$ and $1\mathbf{x} = \mathbf{x}$, $\forall \mathbf{x} \in V$
 - ▶ **Inverse**: $\forall \mathbf{x} \in V, \exists -\mathbf{x} \in V$ such that $\mathbf{x} + (-\mathbf{x}) = \mathbf{0}$
 - ▶ **Distributivity**: $a(\mathbf{x} + \mathbf{y}) = a\mathbf{x} + a\mathbf{y}$ and $(a + b)\mathbf{x} = a\mathbf{x} + b\mathbf{x}$, $\forall a, b \in F$ and $\forall \mathbf{x}, \mathbf{y} \in V$
- ▶ **Examples**: real vectors \mathbb{R}^d , complex vectors \mathbb{C}^d , rational vectors \mathbb{Q}^d , functions $\mathbb{R}^d \mapsto \mathbb{R}$

Basis and Dimension

- ▶ A **basis** of a vector space V over a field F is a set $B \subseteq V$ that satisfies:
 - ▶ **linear independence**: for all finite $\{\mathbf{x}_1, \dots, \mathbf{x}_m\} \subseteq B$,
if $a_1\mathbf{x}_1 + \dots + a_m\mathbf{x}_m = 0$ for some $a_1, \dots, a_m \in F$, then $a_1 = \dots = a_m = 0$
 - ▶ B **spans** V : $\forall \mathbf{x} \in V, \exists \mathbf{x}_1, \dots, \mathbf{x}_d \in B$ and unique $a_1, \dots, a_d \in F$ such
that $\mathbf{x} = a_1\mathbf{x}_1 + \dots + a_d\mathbf{x}_d$
- ▶ The **dimension** d of a vector space V is the cardinality of its bases

Inner Product and Norm

- ▶ An **inner product** on a vector space V over a field F is a function $\langle \cdot, \cdot \rangle : V \times V \mapsto F$ such that for all $a \in F$ and all $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$:
 - ▶ $\langle a\mathbf{x}, \mathbf{y} \rangle = a\langle \mathbf{x}, \mathbf{y} \rangle$ (homogeneity)
 - ▶ $\langle \mathbf{x} + \mathbf{y}, \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{z} \rangle + \langle \mathbf{y}, \mathbf{z} \rangle$ (additivity)
 - ▶ $\langle \mathbf{x}, \mathbf{y} \rangle = \overline{\langle \mathbf{y}, \mathbf{x} \rangle}$ (conjugate symmetry)
 - ▶ $\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$ (non-negativity)
 - ▶ $\langle \mathbf{x}, \mathbf{x} \rangle = 0$ iff $\mathbf{x} = \mathbf{0}$ (definiteness)
- ▶ A **norm** on a vector space V over a field F is a function $\| \cdot \| : V \rightarrow \mathbb{R}$ such that for all $a \in F$ and all $\mathbf{x}, \mathbf{y} \in V$:
 - ▶ $\| a\mathbf{x} \| = |a| \| \mathbf{x} \|$ (absolute homogeneity)
 - ▶ $\| \mathbf{x} + \mathbf{y} \| \leq \| \mathbf{x} \| + \| \mathbf{y} \|$ (triangle inequality)
 - ▶ $\| \mathbf{x} \| \geq 0$ (non-negativity)
 - ▶ $\| \mathbf{x} \| = 0$ iff $\mathbf{x} = \mathbf{0}$ (definiteness)

Euclidean Vector Space

- ▶ A **Euclidean vector space** \mathbb{R}^d is a vector space with finite dimension d over the real numbers \mathbb{R}
- ▶ A **Euclidean vector** $\mathbf{x} \in \mathbb{R}^d$ is a collection of scalars $x_i \in \mathbb{R}$ for $i = 1, \dots, d$ organized as a column:

$$\mathbf{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_d \end{bmatrix}$$

- ▶ The **transpose** of $\mathbf{x} \in \mathbb{R}^d$ is organized as a row: $\mathbf{x}^T = [x_1 \ \cdots \ x_d]$
- ▶ The **Euclidean inner product** between two vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$ is:

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \mathbf{y} = \sum_{i=1}^d x_i y_i$$

- ▶ The **Euclidean norm** of a vector $\mathbf{x} \in \mathbb{R}^d$ is $\|\mathbf{x}\|_2 := \sqrt{\mathbf{x}^T \mathbf{x}}$ and satisfies:
 - ▶ $\max_{1 \leq i \leq d} |x_i| \leq \|\mathbf{x}\|_2 \leq \sqrt{d} \max_{1 \leq i \leq d} |x_i|$
 - ▶ $|\mathbf{x}^T \mathbf{y}| \leq \|\mathbf{x}\|_2 \|\mathbf{y}\|_2$ (Cauchy-Schwarz Inequality)

Matrices

- ▶ A **matrix** $A \in \mathbb{R}^{m \times n}$ is a rectangular array of scalars $A_{ij} \in \mathbb{R}$ for $i = 1, \dots, m$ and $j = 1, \dots, n$
- ▶ The entries of the **transpose** $A^T \in \mathbb{R}^{n \times m}$ of a matrix $A \in \mathbb{R}^{m \times n}$ are $A_{ij}^T = A_{ji}$. The transpose satisfies: $(AB)^T = B^T A^T$
- ▶ The **trace** of a matrix $A \in \mathbb{R}^{n \times n}$ is the sum of its diagonal entries:

$$\text{tr}(A) := \sum_{i=1}^n A_{ii} \qquad \text{tr}(ABC) = \text{tr}(BCA) = \text{tr}(CAB)$$

- ▶ The **Frobenius inner product** between two matrices $X, Y \in \mathbb{R}^{m \times n}$ is:

$$\langle X, Y \rangle = \text{tr}(X^T Y)$$

- ▶ The **Frobenius norm** of a matrix $X \in \mathbb{R}^{m \times n}$ is: $\|X\|_F := \sqrt{\text{tr}(X^T X)}$

Matrix Determinant and Inverse

- ▶ The **determinant** of a matrix $A \in \mathbb{R}^{n \times n}$ is:

$$\det(A) := \sum_{j=1}^n A_{ij} \mathbf{cof}_{ij}(A) \qquad \det(AB) = \det(A) \det(B) = \det(BA)$$

where $\mathbf{cof}_{ij}(A)$ is the **cofactor** of the entry A_{ij} and is equal to $(-1)^{i+j}$ times the determinant of the $(n-1) \times (n-1)$ submatrix that results when the i^{th} -row and j^{th} -col of A are removed. This recursive definition uses the fact that the determinant of a scalar is the scalar itself.

- ▶ The **adjugate** is the transpose of the cofactor matrix:

$$\mathbf{adj}(A) := \mathbf{cof}(A)^{\top}$$

- ▶ The **inverse** A^{-1} of A exists iff $\det(A) \neq 0$ and satisfies:

$$A^{-1} = \frac{\mathbf{adj}(A)}{\det(A)} \qquad (AB)^{-1} = B^{-1}A^{-1}$$

Matrix Inversion Lemma

► **Square completion:**

$$\frac{1}{2}x^\top Ax + b^\top x + c = \frac{1}{2}(x + A^{-1}b)^\top A(x + A^{-1}b) + c - \frac{1}{2}b^\top A^{-1}b$$

► **Woodbury matrix identity:**

$$(A + BDC)^{-1} = A^{-1} - A^{-1}B(CA^{-1}B + D^{-1})^{-1}CA^{-1}$$

► **Block matrix inversion:**

$$\begin{aligned} \begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} &= \begin{bmatrix} I & 0 \\ D^{-1}C & I \end{bmatrix}^{-1} \begin{bmatrix} A - BD^{-1}C & 0 \\ 0 & D \end{bmatrix}^{-1} \begin{bmatrix} I & BD^{-1} \\ 0 & I \end{bmatrix}^{-1} \\ &= \begin{bmatrix} I & 0 \\ -D^{-1}C & I \end{bmatrix} \begin{bmatrix} (A - BD^{-1}C)^{-1} & 0 \\ 0 & D^{-1} \end{bmatrix} \begin{bmatrix} I & -BD^{-1} \\ 0 & I \end{bmatrix} \\ &= \begin{bmatrix} (A - BD^{-1}C)^{-1} & -(A - BD^{-1}C)^{-1}BD^{-1} \\ -D^{-1}C(A - BD^{-1}C)^{-1} & D^{-1} + D^{-1}C(A - BD^{-1}C)^{-1}BD^{-1} \end{bmatrix} \end{aligned}$$

Eigenvalue Decomposition

- ▶ For any $A \in \mathbb{R}^{n \times n}$, if there exists $\mathbf{q} \in \mathbb{C}^n \setminus \{\mathbf{0}\}$ and $\lambda \in \mathbb{C}$ such that:

$$A\mathbf{q} = \lambda\mathbf{q}$$

then \mathbf{q} is an **eigenvector** corresponding to the **eigenvalue** λ .

- ▶ A real matrix can have complex eigenvalues and eigenvectors, which appear in conjugate pairs.
- ▶ Eigenvectors are not unique since for any $c \in \mathbb{C} \setminus \{0\}$, $c\mathbf{q}$ is an eigenvector corresponding to the same eigenvalue.
- ▶ The n eigenvalues of $A \in \mathbb{R}^{n \times n}$ are precisely the n roots of the **characteristic polynomial** of A :

$$p(\lambda) := \det(\lambda I - A)$$

Eigenvalue Decomposition

- ▶ **Diagonalizable matrix:** n linearly independent eigenvectors \mathbf{q}_i can be found for $A \in \mathbb{R}^{n \times n}$: $A\mathbf{q}_i = \lambda_i\mathbf{q}_i$ for $i = 1, \dots, n$
- ▶ If the eigenvalues λ_i of A are distinct, then A is diagonalizable
- ▶ **Eigen decomposition:** if A is diagonalizable, we can stack all n equations $A\mathbf{q}_i = \lambda_i\mathbf{q}_i$ to obtain an eigen decomposition of A :

$$A = Q\Lambda Q^{-1}$$

- ▶ **Jordan decomposition:** any A can be decomposed using an invertible matrix Q of generalized eigenvectors and an upper-triangular matrix J :

$$A = QJQ^{-1}$$

- ▶ **Jordan form J of A :** an upper-triangular block-diagonal matrix:

$$J = \text{diag}(B(\lambda_1, m_1), \dots, B(\lambda_k, m_k))$$

where $\lambda_1, \dots, \lambda_k$ are the eigenvalues of A and $m_1 + \dots + m_k = n$.

$$B(\lambda, m) = \begin{bmatrix} \lambda & 1 & 0 & 0 \\ 0 & \ddots & \ddots & 0 \\ \vdots & \ddots & \lambda & 1 \\ 0 & 0 & 0 & \lambda \end{bmatrix}_{11}$$

Eigenvalue Decomposition

- ▶ The roots of a polynomial are continuous functions of its coefficients and hence the eigenvalues of a matrix are continuous functions of its entries.

$$\operatorname{tr}(A) := \sum_{i=1}^n \lambda_i \qquad \det(A) := \prod_{i=1}^n \lambda_i$$

- ▶ A^\top has the same eigenvalues and eigenvectors as A
- ▶ $A^\top A$ has the same eigenvectors as A but its eigenvalues are λ^2
- ▶ A^k for $k = 1, 2, \dots$ has the same eigenvectors as A but its eigenvalues are λ^k
- ▶ A^{-1} has the same eigenvectors as A but its eigenvalues are λ^{-1}
- ▶ The eigenvalues of A are invariant under any unitary transformation U^*AU for $U^*U = UU^* = I$
- ▶ If A is symmetric, $A^\top = A$, then all its eigenvalues are real and all its eigenvectors may be chosen orthogonal: $Q^{-1} = Q^\top$

Singular Value Decomposition

- ▶ An eigen-decomposition does not exist for $A \in \mathbb{R}^{m \times n}$
- ▶ $A \in \mathbb{R}^{m \times n}$ with rank $r \leq \min\{m, n\}$ can be diagonalized by two orthogonal matrices $U \in \mathbb{R}^{m \times m}$ and $V \in \mathbb{R}^{n \times n}$ via **singular value decomposition**:

$$A = U\Sigma V^T \quad \Sigma = \begin{bmatrix} \sigma_1 & & & \\ & \ddots & & \\ & & \sigma_r & \\ & & & \end{bmatrix} \in \mathbb{R}^{m \times n}$$

- ▶ U contains the m orthogonal eigenvectors of the symmetric matrix $AA^T \in \mathbb{R}^{m \times m}$ and satisfies $U^T U = U U^T = I$
- ▶ V contains the n orthogonal eigenvectors of the symmetric matrix $A^T A \in \mathbb{R}^{n \times n}$ and satisfies $V^T V = V V^T = I$
- ▶ Σ contains the singular values $\sigma_i = \sqrt{\lambda_i}$, equal to the square roots of the r non-zero eigenvalues λ_i of AA^T or $A^T A$, on its diagonal
- ▶ If A is normal ($A^T A = A A^T$), its singular values are related to its eigenvalues via $\sigma_i = |\lambda_i|$

Matrix Pseudo Inverse

- ▶ The **pseudo-inverse** $A^\dagger \in \mathbb{R}^{n \times m}$ of $A \in \mathbb{R}^{m \times n}$ can be obtained from its SVD $A = U\Sigma V^T$:

$$A^\dagger = V\Sigma^\dagger U^T \quad \Sigma^\dagger = \begin{bmatrix} 1/\sigma_1 & & & \\ & \ddots & & \\ & & 1/\sigma_r & \\ & & & \end{bmatrix} \in \mathbb{R}^{n \times m}$$

- ▶ The pseudo-inverse $A^\dagger \in \mathbb{R}^{n \times m}$ satisfies the Moore-Penrose conditions:
 - ▶ $AA^\dagger A = A$
 - ▶ $A^\dagger AA^\dagger = A^\dagger$
 - ▶ $(AA^\dagger)^T = AA^\dagger$
 - ▶ $(A^\dagger A)^T = A^\dagger A$

Linear System of Equations

- ▶ Consider the linear system of equations $A\mathbf{x} = \mathbf{b}$ for $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{b} \in \mathbb{R}^m$, and $A \in \mathbb{R}^{m \times n}$ with SVD $A = U\Sigma V^T$ and rank r
- ▶ The **column space** or **image** of A is $im(A) \subseteq \mathbb{R}^m$ and is spanned by the r columns of U corresponding to non-zero singular values
- ▶ The **null space** or **kernel** of A is $ker(A) \subseteq \mathbb{R}^n$ and is spanned by the $n - r$ columns of V corresponding to zero singular values
- ▶ The **row space** or **co-image** of A is $im(A^T) \subseteq \mathbb{R}^n$ and is spanned by the r columns of V corresponding to non-zero singular values
- ▶ The **left null space** or **co-kernel** of A is $ker(A^T) \subseteq \mathbb{R}^m$ and is spanned by the $m - r$ columns of U corresponding to zero singular values
- ▶ The **domain** of A is $\mathbb{R}^n = ker(A) \oplus im(A^T)$
- ▶ The **co-domain** of A is $\mathbb{R}^m = ker(A^T) \oplus im(A)$

Solution of Linear System of Equations

- ▶ Consider the linear system of equations $A\mathbf{x} = \mathbf{b}$ for $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{b} \in \mathbb{R}^m$, and $A \in \mathbb{R}^{m \times n}$ with SVD $A = U\Sigma V^T$ and rank r
- ▶ If $\mathbf{b} \in im(A)$, i.e., $\mathbf{b}^T \mathbf{v} = 0$ for all $\mathbf{v} \in ker(A^T)$, then $A\mathbf{x} = \mathbf{b}$ has **one or infinitely many solutions** $\mathbf{x} = A^\dagger \mathbf{b} + (I - A^\dagger A)\mathbf{y}$ for any $\mathbf{y} \in \mathbb{R}^n$
- ▶ If $\mathbf{b} \notin im(A)$, then **no solution exists** and $\mathbf{x} = A^\dagger \mathbf{b}$ is an approximate solution with minimum $\|\mathbf{x}\|$ and $\|A\mathbf{x} - \mathbf{b}\|$ norms
- ▶ If $m = n = r$, then $A\mathbf{x} = \mathbf{b}$ has a **unique solution** $\mathbf{x} = A^\dagger \mathbf{b} = A^{-1}\mathbf{b}$

Positive Semidefinite Matrices

- ▶ The product $\mathbf{x}^\top A \mathbf{x}$ for $A \in \mathbb{R}^{n \times n}$ and $\mathbf{x} \in \mathbb{R}^n$ is called a **quadratic form** and A can be assumed **symmetric**, $A = A^\top$, because:

$$\frac{1}{2} \mathbf{x}^\top (A + A^\top) \mathbf{x} = \mathbf{x}^\top A \mathbf{x}, \quad \forall \mathbf{x} \in \mathbb{R}^n$$

- ▶ A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is **positive semidefinite** if $\mathbf{x}^\top A \mathbf{x} \geq 0$ for all $\mathbf{x} \in \mathbb{R}^n$.
- ▶ A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is **positive definite** if it is positive semidefinite and if $\mathbf{x}^\top A \mathbf{x} = 0$ implies $\mathbf{x} = 0$.
- ▶ All eigenvalues of a symmetric positive semidefinite matrix are non-negative.
- ▶ All eigenvalues of a symmetric positive definite matrix are positive.

Schur Complement

- ▶ The Schur complement of block D of $M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$ is $S_D = A - BD^{-1}C$
- ▶ The Schur complement of block A of $M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$ is $S_A = D - CA^{-1}B$
- ▶ Let $M = \begin{bmatrix} A & B \\ B^\top & D \end{bmatrix}$ be symmetric. Then:
 - ▶ $M \succ 0 \Leftrightarrow A \succ 0, S_A = D - B^\top A^{-1}B \succ 0$
 - ▶ $M \succ 0 \Leftrightarrow D \succ 0, S_D = A - BD^{-1}B^\top \succ 0$
 - ▶ $M \succeq 0 \Leftrightarrow A \succeq 0, S_A \succeq 0, (I - AA^\dagger)B = 0$
 - ▶ $M \succeq 0 \Leftrightarrow D \succeq 0, S_D \succeq 0, (I - DD^\dagger)B^\top = 0$

Derivatives (numerator layout)

- Derivatives of $y \in \mathbb{R}^m$ and $Y \in \mathbb{R}^{m \times n}$ by scalar $x \in \mathbb{R}$:

$$\frac{dy}{dx} = \begin{bmatrix} \frac{dy_1}{dx} \\ \vdots \\ \frac{dy_m}{dx} \end{bmatrix} \in \mathbb{R}^{m \times 1} \quad \frac{dY}{dx} = \begin{bmatrix} \frac{dY_{11}}{dx} & \cdots & \frac{dY_{1n}}{dx} \\ \vdots & \ddots & \vdots \\ \frac{dY_{m1}}{dx} & \cdots & \frac{dY_{mn}}{dx} \end{bmatrix} \in \mathbb{R}^{m \times n}$$

- Derivatives of $y \in \mathbb{R}$ and $\mathbf{y} \in \mathbb{R}^m$ by vector $\mathbf{x} \in \mathbb{R}^p$:

$$\frac{dy}{d\mathbf{x}} = \underbrace{\begin{bmatrix} \frac{dy}{dx_1} & \cdots & \frac{dy}{dx_p} \end{bmatrix}}_{[\nabla_{\mathbf{x}} y]^T \text{ (gradient transpose)}} \in \mathbb{R}^{1 \times p} \quad \frac{d\mathbf{y}}{d\mathbf{x}} = \underbrace{\begin{bmatrix} \frac{dy_1}{dx_1} & \cdots & \frac{dy_1}{dx_p} \\ \vdots & \ddots & \vdots \\ \frac{dy_m}{dx_1} & \cdots & \frac{dy_m}{dx_p} \end{bmatrix}}_{\text{Jacobian}} \in \mathbb{R}^{m \times p}$$

- Derivative of $y \in \mathbb{R}$ by matrix $X \in \mathbb{R}^{p \times q}$:

$$\frac{dy}{dX} = \begin{bmatrix} \frac{dy}{dX_{11}} & \cdots & \frac{dy}{dX_{p1}} \\ \vdots & \ddots & \vdots \\ \frac{dy}{dX_{1q}} & \cdots & \frac{dy}{dX_{pq}} \end{bmatrix} \in \mathbb{R}^{q \times p}$$

Matrix Derivatives Example

- ▶ $\frac{d}{dX_{ij}} X = \mathbf{e}_i \mathbf{e}_j^\top$
- ▶ $\frac{d}{dx} A\mathbf{x} = A$
- ▶ $\frac{d}{dx} \mathbf{x}^\top A\mathbf{x} = \mathbf{x}^\top (A + A^\top)$
- ▶ $\frac{d}{dx} M^{-1}(x) = -M^{-1}(x) \frac{dM(x)}{dx} M^{-1}(x)$
- ▶ $\frac{d}{dX} \text{tr}(AX^{-1}B) = -X^{-1}BAX^{-1}$
- ▶ $\frac{d}{dX} \log \det X = X^{-1}$

Matrix Derivatives Example

$$\blacktriangleright \frac{d}{dx} \mathbf{Ax} = \begin{bmatrix} \frac{d}{dx_1} \sum_{j=1}^n A_{1j}x_j & \cdots & \frac{d}{dx_n} \sum_{j=1}^n A_{1j}x_j \\ \vdots & \ddots & \vdots \\ \frac{d}{dx_1} \sum_{j=1}^n A_{mj}x_j & \cdots & \frac{d}{dx_n} \sum_{j=1}^n A_{mj}x_j \end{bmatrix} = \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & \ddots & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix}$$

$$\blacktriangleright \frac{d}{dx} \mathbf{x}^\top \mathbf{Ax} = \mathbf{x}^\top \mathbf{A}^\top \frac{d\mathbf{x}}{dx} + \mathbf{x}^\top \frac{d\mathbf{Ax}}{dx} = \mathbf{x}^\top (\mathbf{A}^\top + \mathbf{A})$$

$$\blacktriangleright M(x)M^{-1}(x) = \mathbf{I} \Rightarrow 0 = \left[\frac{d}{dx} M(x) \right] M^{-1}(x) + M(x) \left[\frac{d}{dx} M^{-1}(x) \right]$$

$$\begin{aligned} \blacktriangleright \frac{d}{dX_{ij}} \text{tr}(\mathbf{AX}^{-1}\mathbf{B}) &= \text{tr}\left(\mathbf{A} \frac{d}{dX_{ij}} \mathbf{X}^{-1} \mathbf{B}\right) = -\text{tr}(\mathbf{AX}^{-1} \mathbf{e}_i \mathbf{e}_j^\top \mathbf{X}^{-1} \mathbf{B}) \\ &= -\mathbf{e}_j^\top \mathbf{X}^{-1} \mathbf{B} \mathbf{A} \mathbf{X}^{-1} \mathbf{e}_i = -\mathbf{e}_i^\top (\mathbf{X}^{-1} \mathbf{B} \mathbf{A} \mathbf{X}^{-1})^\top \mathbf{e}_j \end{aligned}$$

$$\begin{aligned} \blacktriangleright \frac{d}{dX_{ij}} \log \det X &= \frac{1}{\det(X)} \frac{d}{dX_{ij}} \sum_{k=1}^n X_{ik} \mathbf{cof}_{ik}(X) \\ &= \frac{1}{\det(X)} \mathbf{cof}_{ij}(X) = \frac{1}{\det(X)} \mathbf{adj}_{ji}(X) = \mathbf{e}_i^\top \mathbf{X}^{-T} \mathbf{e}_j \end{aligned}$$