

ECE276B: Planning & Learning in Robotics

Lecture 2: Markov Chains

Nikolay Atanasov
natanasov@ucsd.edu

UC San Diego
JACOBS SCHOOL OF ENGINEERING
Electrical and Computer Engineering

Outline

Markov Chains

Absorbing Markov Chains

Ergodic Markov Chains

Markov Chain

- ▶ **Stochastic process:** indexed collection of random variables $\{x_0, x_1, \dots\}$
- ▶ **Markov chain:** memoryless stochastic process $\{x_0, x_1, \dots\}$:
 - ▶ x_0 has probability density function $p_0(\cdot)$
 - ▶ x_{t+1} conditioned on x_t has probability density function $p_f(\cdot | x_t)$ and is independent of the history $x_{0:t-1}$
- ▶ **Markov assumption:**
"The future is independent of the past given the present"


Markov Chain

Stochastic process defined by a tuple (\mathcal{X}, p_0, p_f) :

- ▶ \mathcal{X} is a discrete or continuous space
- ▶ $p_0(\cdot)$ is a prior pdf defined on \mathcal{X}
- ▶ $p_f(\cdot | \mathbf{x})$ is a conditional pdf defined on \mathcal{X} for given $\mathbf{x} \in \mathcal{X}$ that specifies the stochastic process transitions
- ▶ When the state space is finite, $\mathcal{X} := \{1, \dots, N\}$, the pdf p_f can be represented by an $N \times N$ transition matrix with elements:

$$P_{ij} := \mathbb{P}(x_{t+1} = j | x_t = i) = p_f(j | x_t = i)$$


Example: Student Markov Chain



Example: Student Markov Chain

► Sample paths:

- C1 C2 C3 Pass Sleep
- C1 FB FB C1 C2 Sleep
- C1 C2 C3 Pub C2 C3 Pass Sleep
- C1 FB FB C1 C2 C3 Pub C1 FB FB
FB C1 C2 Sleep



► Transition matrix:

$$P = \begin{matrix} FB \\ C1 \\ C2 \\ C3 \\ Pub \\ Pass \\ Sleep \end{matrix} \begin{bmatrix} 0.9 & 0.1 & 0 & 0 & 0 & 0 & 0 \\ 0.5 & 0 & 0.5 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.8 & 0 & 0 & 0.2 \\ 0 & 0 & 0 & 0 & 0.4 & 0.6 & 0 \\ 0 & 0.2 & 0.4 & 0.4 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Chapman-Kolmogorov Equation

- ▶ n -step transition probabilities of Markov chain on $\mathcal{X} = \{1, \dots, N\}$

$$P_{ij}^{(n)} := \mathbb{P}(x_{t+n} = j \mid x_t = i) = \mathbb{P}(x_n = j \mid x_0 = i)$$

- ▶ **Chapman-Kolmogorov:** the n -step transition probabilities can be obtained recursively from the 1-step transition probabilities:

$$P_{ij}^{(n)} = \sum_{k=1}^N P_{ik}^{(m)} P_{kj}^{(n-m)}, \quad \forall i, j, n, 0 \leq m \leq n$$
$$P^{(n)} = \underbrace{P \dots P}_{n \text{ times}} = P^n$$

- ▶ Given the transition matrix P and a vector $\mathbf{p}_0 := [p_0(1), \dots, p_0(N)]^\top$ of prior probabilities, the vector of probabilities \mathbf{p}_n after n steps is:

$$\mathbf{p}_n^\top = \mathbf{p}_0^\top P^n$$

Example: Student Markov Chain

$$P = \begin{matrix} FB \\ C1 \\ C2 \\ C3 \\ Pub \\ Pass \\ Sleep \end{matrix} \begin{bmatrix} 0.9 & 0.1 & 0 & 0 & 0 & 0 & 0 \\ 0.5 & 0 & 0.5 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.8 & 0 & 0 & 0.2 \\ 0 & 0 & 0 & 0 & 0.4 & 0.6 & 0 \\ 0 & 0.2 & 0.4 & 0.4 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$P^2 = \begin{matrix} FB \\ C1 \\ C2 \\ C3 \\ Pub \\ Pass \\ Sleep \end{matrix} \begin{bmatrix} 0.86 & 0.09 & 0.05 & 0 & 0 & 0 & 0 \\ 0.45 & 0.05 & 0 & 0.4 & 0 & 0 & 0.1 \\ 0 & 0 & 0 & 0 & 0.32 & 0.48 & 0.2 \\ 0 & 0.08 & 0.16 & 0.16 & 0 & 0 & 0.6 \\ 0.1 & 0 & 0.1 & 0.32 & 0.16 & 0.24 & 0.08 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$P^{100} = \begin{matrix} FB \\ C1 \\ C2 \\ C3 \\ Pub \\ Pass \\ Sleep \end{matrix} \begin{bmatrix} 0.01 & 0 & 0 & 0 & 0 & 0 & 0.99 \\ 0.01 & 0 & 0 & 0 & 0 & 0 & 0.99 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

First Passage Time

- ▶ **First passage time:** the number of transitions necessary to reach state j for the first time is a random variable:

$$\tau_j := \min\{t \geq 1 \mid x_t = j\}$$

- ▶ **Recurrence time:** the first passage time τ_i from $x_0 = i$ to $j = i$
- ▶ **Probability of first passage in n steps:** $\rho_{ij}^{(n)} := \mathbb{P}(\tau_j = n \mid x_0 = i)$

$$\rho_{ij}^{(1)} = P_{ij}$$

$$\rho_{ij}^{(2)} = [P^2]_{ij} - \rho_{ij}^{(1)} P_{jj} \quad (\text{first time we visit } j \text{ should not be } 1!)$$

⋮

$$\rho_{ij}^{(n)} = [P^n]_{ij} - \rho_{ij}^{(1)} [P^{n-1}]_{jj} - \rho_{ij}^{(2)} [P^{n-2}]_{jj} - \dots - \rho_{ij}^{(n-1)} P_{jj}$$

- ▶ **Probability of first passage:** $\rho_{ij} := \mathbb{P}(\tau_j < \infty \mid x_0 = i) = \sum_{n=1}^{\infty} \rho_{ij}^{(n)}$
- ▶ **Number of visits to j up to time n :**

$$v_j^{(n)} := \sum_{t=0}^n \mathbb{1}\{x_t = j\} \quad v_j := \lim_{n \rightarrow \infty} v_j^{(n)}$$

Recurrence and Transience

- ▶ **Absorbing state:** a state j such that $P_{jj} = 1$
- ▶ **Transient state:** a state j such that $\rho_{jj} < 1$
- ▶ **Recurrent state:** a state j such that $\rho_{jj} = 1$
- ▶ **Positive recurrent state:** a recurrent state j with $\mathbb{E}[\tau_j \mid x_0 = j] < \infty$
- ▶ **Null recurrent state:** a recurrent state j with $\mathbb{E}[\tau_j \mid x_0 = j] = \infty$
- ▶ **Periodic state:** can only be visited at integer multiples of t
- ▶ **Ergodic state:** a positive recurrent state that is aperiodic

Recurrence and Transience

Total Number of Visits Lemma

$$\mathbb{P}(v_j \geq k + 1 \mid x_0 = j) = \rho_{jj}^k \text{ for all } k \geq 0$$

Proof:

By Markov property and induction: $\mathbb{P}(v_j \geq k + 1 \mid x_0 = j) = \rho_{jj} \mathbb{P}(v_j \geq k \mid x_0 = j)$.

0-1 Law for the Total Number of Visits

$$j \text{ is recurrent iff } \mathbb{E}[v_j \mid x_0 = j] = \infty$$

Proof: Since v_j is discrete, we can write $v_j = \sum_{k=0}^{\infty} \mathbb{1}\{v_j > k\}$ and

$$\mathbb{E}[v_j \mid x_0 = j] = \sum_{k=0}^{\infty} \mathbb{P}(v_j \geq k + 1 \mid x_0 = j) = \sum_{k=0}^{\infty} \rho_{jj}^k = \frac{\rho_{jj}}{1 - \rho_{jj}}$$

Recurrence Is Contagious

$$i \text{ is recurrent and } \rho_{ij} > 0 \quad \Rightarrow \quad j \text{ is recurrent and } \rho_{ji} = 1$$

Mean First Passage Time

▶ **Mean first passage time:** $M_{ij} := \mathbb{E}[\tau_j \mid x_0 = i]$

▶ By the law of total probability:

$$M_{ij} = P_{ij} + \sum_{k \neq j} P_{ik}(1 + M_{kj}) = 1 + \sum_{k \neq j} P_{ik}M_{kj}$$

▶ Let $M \in \mathbb{R}^{N \times N}$ with elements M_{ij} contain all mean first passage times


▶ The matrix of mean first passage times satisfies:

$$M = \mathbf{1}\mathbf{1}^\top + P(M - D)$$

where $D = \mathbf{diag}(M_{11}, \dots, M_{NN})$

Equivalence Classes

- ▶ $i \rightarrow j$: state j is **accessible** from state i if $P_{ij}^{(n)} > 0$ for some n
- ▶ Every state is accessible from itself since $P_{ii}^{(0)} = 1$
- ▶ $i \leftrightarrow j$: state i and j **communicate** if they are accessible from each other
- ▶ **Equivalence class**: a set of states which communicate with each other
- ▶ **Example**: find the equivalence classes for this Markov chain



Classification of Markov Chains

- ▶ **Absorbing Markov Chain:** contains at least one absorbing state that can be reached from every other state (not necessarily in one step)
- ▶ **Irreducible Markov Chain:** all states communicate with each other
- ▶ **Ergodic Markov Chain:** an aperiodic, irreducible and positive recurrent Markov chain

Periodicity

- ▶ Periodicity has important role in the long-term behavior of a Markov chain
- ▶ The **period** of a state i is the largest integer d_i such that $P_{ii}^{(n)} = 0$ whenever n is not divisible by d_i
 - ▶ If $d_i > 1$, then i is **periodic**
 - ▶ If $d_i = 1$, then i is **aperiodic**
- ▶ If $i \leftrightarrow j$, then $d_i = d_j$. Hence, all states of an irreducible Markov chain have the same period.
- ▶ Two integers are **co-prime** if their greatest common divisor (gcd) is 1
- ▶ If we can find co-prime l and m such that $P_{ii}^{(l)} > 0$ and $P_{ii}^{(m)} > 0$, then i is aperiodic
- ▶ Since 1 is co-prime to every integer, any state i with a self-transition is aperiodic

Periodicity

- ▶ A matrix P is **non-negative** if all $P_{ij} \geq 0$
- ▶ A matrix P is **stochastic** if its rows sum to 1, i.e., $\sum_j P_{ij} = 1$ for all i
- ▶ A non-negative matrix P is **quasi-positive** if there exists a natural number $m \geq 1$ such that all entries of P^m are strictly positive
- ▶ If P is a stochastic matrix and is quasi-positive, i.e., all entries of P^m are positive, then for all $n \geq m$ all entries of P^n are positive
- ▶ **Aperiodicity Lemma:** A stochastic transition matrix P is irreducible and aperiodic if and only if P is **quasi-positive**.
- ▶ A finite Markov chain with transition matrix P is ergodic if and only if P is **quasi-positive**

Stationary and Limiting Distributions

▶ **Stationary distribution:** a vector $\mathbf{w} \in \{\mathbf{p} \in [0, 1]^N \mid \mathbf{1}^\top \mathbf{p} = 1\}$ such that $\mathbf{w}^\top P = \mathbf{w}^\top$

▶ **Limiting distribution:** a vector $\mathbf{w} \in \{\mathbf{p} \in [0, 1]^N \mid \mathbf{1}^\top \mathbf{p} = 1\}$ such that:

$$\lim_{t \rightarrow \infty} \mathbb{P}(x_t = j \mid x_0 = i) = \mathbf{w}_j$$

▶ If it exists, the limiting distribution of a Markov chain is stationary

▶ **Absorbing chains** have limiting distributions with nonzero elements only in absorbing states

▶ **Ergodic chains** have a unique limiting distribution (Perron-Frobenius Thm)

▶ **Periodic chains** may not have a limiting distribution; their stationary distribution has $w_j > 0$ only for recurrent states and w_j is the frequency $\frac{v_j^{(n)}}{n+1}$ of being in state j as $n \rightarrow \infty$

Example

- ▶ Consider a Markov chain with:
 - ▶ state space $\mathcal{X} = \{0, 1\}$
 - ▶ prior pmf $\mathbf{p}_0 = [\mathbb{P}(x_0 = 0), \mathbb{P}(x_0 = 1)]^\top = [\gamma, 1 - \gamma]^\top$
 - ▶ transition matrix with $a, b \in [0, 1]$, $0 < a + b < 2$:

$$P = \begin{bmatrix} 1 - a & a \\ b & 1 - b \end{bmatrix}$$

- ▶ By induction: $P^n = \frac{1}{a+b} \begin{bmatrix} b & a \\ b & a \end{bmatrix} + \frac{(1-a-b)^n}{a+b} \begin{bmatrix} a & -a \\ -b & b \end{bmatrix}$
- ▶ Since $-1 < 1 - a - b < 1$: $\lim_{n \rightarrow \infty} P^n = \frac{1}{a+b} \begin{bmatrix} b & a \\ b & a \end{bmatrix}$
- ▶ Limiting distribution: exists and is not dependent on the initial pmf \mathbf{p}_0 :

$$\lim_{t \rightarrow \infty} \mathbf{p}_t^\top = \lim_{t \rightarrow \infty} \mathbf{p}_0^\top P^t = \frac{1}{a+b} \mathbf{p}_0^\top \begin{bmatrix} b & a \\ b & a \end{bmatrix} = \left[\frac{b}{a+b}, \frac{b}{a+b} \right]$$

Example

▶ If $a = b = 1$, the transition matrix is $P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

▶ This Markov chain is periodic:

$$x_t = \begin{cases} x_0 & \text{if } t \text{ is even} \\ x_1 & \text{if } t \text{ is odd} \end{cases}$$

▶ Stationary distribution: $\mathbf{w} = [\frac{1}{2}, \frac{1}{2}]$

▶ Limiting distribution: does not exist. The pmf \mathbf{p}_t does not converge as $t \rightarrow \infty$ and depends on \mathbf{p}_0

Outline

Markov Chains

Absorbing Markov Chains

Ergodic Markov Chains

Absorbing Markov Chains

► Interesting questions:

Q1: On average, how many times is the process in state j ?

Q2: What is the probability that the state will eventually be absorbed?

Q3: What is the expected absorption time?

Q4: What is the probability of being absorbed by j given that we started in i ?

Absorbing Markov Chains

- ▶ **Canonical form:** reorder states so that transient come first: $P = \begin{bmatrix} Q & R \\ 0 & I \end{bmatrix}$

- ▶ One can show that $P^n = \begin{bmatrix} Q^n & * \\ 0 & I \end{bmatrix}$ and $Q^n \rightarrow 0$ as $n \rightarrow \infty$

Proof: If j is transient, then $\rho_{ij} < 1$ and from the 0-1 Law:

$$\infty > \mathbb{E}[v_j \mid x_0 = i] = \mathbb{E}\left[\sum_{n=0}^{\infty} \mathbb{1}\{x_n = j\} \mid x_0 = i\right] = \sum_{n=0}^{\infty} [P^n]_{ij}$$


- ▶ **Fundamental matrix:** $Z^A = (I - Q)^{-1} = \sum_{n=0}^{\infty} Q^n$
 - ▶ Expected number of times the chain is in state j : $Z_{ij}^A = \mathbb{E}[v_j \mid x_0 = i]$
 - ▶ Expected absorption time when starting from state i : $\sum_j Z_{ij}^A$
- ▶ **Absorption probability:** let B_{ij} be the the probability of reaching absorbing state j starting from transient state i :

$$B_{ij} = P_{ij} + \sum_{k \in \text{Transient}} P_{ik} B_{kj} \quad \Rightarrow \quad B = R + QB \quad \Rightarrow \quad B = Z^A R$$

Example: Drunkard's Walk

- ▶ Transition matrix:

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0.5 & 0 & 0.5 & 0 & 0 \\ 0 & 0.5 & 0 & 0.5 & 0 \\ 0 & 0 & 0.5 & 0 & 0.5 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$



- ▶ Canonical form:

$$P = \begin{bmatrix} 0 & 0.5 & 0 & 0.5 & 0 \\ 0.5 & 0 & 0.5 & 0 & 0 \\ 0 & 0.5 & 0 & 0 & 0.5 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

- ▶ Fundamental matrix:

$$Z^A = (I - Q)^{-1} = \begin{bmatrix} 1.5 & 1 & 0.5 \\ 1 & 2 & 1 \\ 0.5 & 1 & 1.5 \end{bmatrix}$$

Outline

Markov Chains

Absorbing Markov Chains

Ergodic Markov Chains

General Finite Markov Chain

- ▶ A finite Markov chain might have several transient and recurrent classes
- ▶ As t increases, the chain is absorbed in one of the recurrent classes
- ▶ We can replace each recurrent class with an absorbing state to obtain a chain with only transient and absorbing states
- ▶ We can obtain the absorption probabilities from $B = Z^A R$
- ▶ Each recurrent class can then be analyzed separately

Perron-Frobenius Theorem (Finite Ergodic Markov Chain)

Theorem

Consider an irreducible, aperiodic, finite Markov chain with transition matrix P . Then, the following hold:

- ▶ 1 is the eigenvalue of max modulus, i.e., $|\lambda| < 1$ for all other eigenvalues
- ▶ 1 is a simple eigenvalue, i.e., the associated eigenspace and left-eigenspace have dimension 1
- ▶ The eigenvector associated with 1 is $\mathbf{1}$
- ▶ The unique left eigenvector \mathbf{w} is nonnegative and $\lim_{n \rightarrow \infty} P^n = \mathbf{1}\mathbf{w}^\top$. Hence, the unique stationary distribution \mathbf{w} is a limiting distribution for the Markov chain, i.e., any initial distribution converges to \mathbf{w} .

Perron-Frobenius Theorem (Ergodic Markov Chain)

Theorem

Consider an irreducible, aperiodic, countably infinite Markov chain. Then, one of the following holds:

- ▶ All states are transient and $\lim_{t \rightarrow \infty} \mathbb{P}(x_t = j | x_0 = i) = 0, \forall i, j$
- ▶ All states are null-recurrent and $\lim_{t \rightarrow \infty} \mathbb{P}(x_t = j | x_0 = i) = 0, \forall i, j$
- ▶ All states are positive-recurrent and there exists a limiting distribution $\mathbf{w}_j = \sum_i \mathbf{w}_i P_{ij}, \sum_j \mathbf{w}_j = 1$ such that:

$$\lim_{t \rightarrow \infty} \mathbb{P}(x_t = j | x_0 = i) = \mathbf{w}_j > 0$$

Fundamental Matrix for Ergodic Chains

- ▶ We can try to define a fundamental matrix as in the absorbing case but $(I - P)^{-1}$ does not exist because $P\mathbf{1} = \mathbf{1}$ (Perron-Frobenius)
- ▶ For absorbing chain, $I + Q + Q^2 + \dots = (I - Q)^{-1}$ converges because $Q^n \rightarrow 0$
- ▶ For ergodic chain, $I + (P - \mathbf{1}\mathbf{w}^\top) + (P^2 - \mathbf{1}\mathbf{w}^\top) + \dots$ converges because $P^n \rightarrow \mathbf{1}\mathbf{w}^\top$ (Perron-Frobenius)
- ▶ Note that $P\mathbf{1}\mathbf{w}^\top = \mathbf{1}\mathbf{w}^\top$ and $(\mathbf{1}\mathbf{w}^\top)^2 = \mathbf{1}\mathbf{w}^\top \mathbf{1}\mathbf{w}^\top = \mathbf{1}\mathbf{w}^\top$

$$\begin{aligned}(P - \mathbf{1}\mathbf{w}^\top)^n &= \sum_{i=0}^n (-1)^i \binom{n}{i} P^{n-i} (\mathbf{1}\mathbf{w}^\top)^i = P^n + \sum_{i=1}^n (-1)^i \binom{n}{i} (\mathbf{1}\mathbf{w}^\top)^i \\ &= P^n + \underbrace{\left[\sum_{i=1}^n (-1)^i \binom{n}{i} \right]}_{(1-1)^{n-1}} (\mathbf{1}\mathbf{w}^\top) = P^n - \mathbf{1}\mathbf{w}^\top\end{aligned}$$

- ▶ Thus, the following inverse exists:

$$I + \sum_{n=1}^{\infty} (P^n - \mathbf{1}\mathbf{w}^\top) = I + \sum_{n=1}^{\infty} (P - \mathbf{1}\mathbf{w}^\top)^n = (I - P + \mathbf{1}\mathbf{w}^\top)^{-1}$$

Fundamental Matrix for Ergodic Chains

- ▶ Consider an ergodic Markov chain with transition matrix P and stationary distribution \mathbf{w}
- ▶ **Fundamental matrix:** $Z^E := (I - P + \mathbf{1}\mathbf{w}^\top)^{-1}$
 - ▶ $\mathbf{w}^\top Z^E = \mathbf{w}^\top$
 - ▶ $Z^E \mathbf{1} = \mathbf{1}$
 - ▶ $Z^E(I - P) = I - \mathbf{1}\mathbf{w}^\top$
- ▶ **Mean first passage time:**
 - ▶ $M_{ij} = \mathbb{E}[\tau_j \mid x_0 = i] = \frac{Z_{jj}^E - Z_{ij}^E}{w_j}, i \neq j$
 - ▶ $M_{ii} = \mathbb{E}[\tau_i \mid x_0 = i] = \frac{1}{w_i}$

Example: Land of Oz

- ▶ Transition matrix:

$$P = \begin{bmatrix} 0.5 & 0.25 & 0.25 \\ 0.5 & 0 & 0.5 \\ 0.25 & 0.25 & 0.5 \end{bmatrix}$$

- ▶ Stationary distribution:

$$\mathbf{w}^\top = [0.4 \quad 0.2 \quad 0.4]$$

- ▶ Fundamental matrix:

$$I - P + \mathbf{1}\mathbf{w}^\top = \begin{bmatrix} 0.9 & -0.05 & 0.15 \\ -0.1 & 1.2 & -0.1 \\ 0.15 & -0.05 & 0.9 \end{bmatrix}$$
$$Z^E = \begin{bmatrix} 1.147 & 0.04 & -0.187 \\ 0.08 & 0.84 & 0.08 \\ -0.187 & 0.04 & 1.147 \end{bmatrix}$$

- ▶ Mean first passage time:

$$M_{12} = \frac{Z_{22}^E - Z_{12}^E}{w_2} = \frac{0.84 - 0.04}{0.2} = 4$$

